

COMMUTATIVE/NONCOMMUTATIVE RANK OF LINEAR MATRICES AND SUBSPACES OF MATRICES OF LOW RANK

MARC FORTIN* AND CHRISTOPHE REUTENAUER†

Dédié à notre ami Alain Lascoux

ABSTRACT. A space of matrix of low rank is a vector space of rectangular matrices whose maximum rank is strictly smaller than the number of rows and the numbers of columns. Among these are the compression spaces, where the rank condition is guaranteed by a rectangular hole of 0's of appropriate size. Spaces of matrices are naturally encoded by linear matrices. The latter have a double existence: over the rational function field, and over the free field (noncommutative). We show that a linear matrix corresponds to a compression space if and only if its rank over both fields is equal. We give a simple linear-algebraic algorithm in order to decide if a given space of matrices is a compression space. We give inequalities relating the commutative rank and the noncommutative rank of a linear matrix.

1. INTRODUCTION

We consider here *linear matrices*, that is, matrices whose entries are of the form $a_0 + a_1x_1 + \cdots + a_dx_d$, where the coefficients a_i are taken in a (commutative) field k and where the x_i are indeterminates, which may be commuting or noncommuting. Such a matrix is of the form $M_0 + \sum_{i=1}^d x_i M_i$, where the M_i are all over k and of the same size.

Linear matrices appear in several area of mathematics: Kronecker pencils ([G] chapter 2), determinantal varieties ([H] chapter 9), spaces of matrices of low rank [EH], algebraic automata theory ([E] VII. 6), free algebras [Ro], free fields [CR], 3-tensors ([K] 4.6.4).

*Marc Fortin; Cégep Régional de Lanaudière à l'Assomption, 180, rue Dorval, l'Assomption (Québec) Canada, J5W 6C1.

†Christophe Reutenauer; Université du Québec à Montréal; Dpartement de mathématiques; Case postale 8888, succursale Centre-Ville, Montréal (Québec) Canada, H3C 3P8 (mailing address); e-mail: christo@math.uqam.ca. Partially supported by NSERC and CRC (Canada).

A linear matrix M has two lives: it may be considered as a matrix over the rational function field $k(x_1, \dots, x_d)$ in the commuting variables x_1, \dots, x_d . It may also be considered as a matrix over the *free field* $k \langle x_1, \dots, x_d \rangle$ (which is the noncommutative analogue of the previous one, see [C1], [C2]). In both cases, M has a rank, which we denote by $crk(M)$ and $ncrk(M)$, respectively. It is intuitively clear that $crk(M) \leq ncrk(M)$ (see Corollary 2.2). One may have strict inequality. For instance, the matrix

$$\begin{pmatrix} 0 & x & y \\ -x & 0 & 1 \\ -y & -1 & 0 \end{pmatrix}$$

has commutative rank 2, but is invertible over the free field, hence of noncommutative rank 3: if one inverts this matrix, one is lead to the element $(yx - xy)^{-1}$, which exists only noncommutatively.

Our main result gives a necessary and sufficient condition for equality of both ranks. To state it, we need some definitions. Observe first that if the commutative rank of M is maximal with respect to its size (that is, equal to $\min(n, p)$ where M is of size $n \times p$), then both ranks coincide, as seen from the previous inequality. So we may disregard this case and may assume that M has rank $< \min(n, p)$. Associate to the linear matrix $M = M_0 + \sum_{i=1}^d x_i M_i$ the subspace H of $k^{n \times p}$ spanned by the matrices M_i .

Since the commutative rank of M is $< \min(n, p)$, the rank of each element of H is also $< \min(n, p)$. Moreover, if we assume that k is *infinite*, the maximum rank in H is equal to the commutative rank of M (see Lemma 3.1). A subspace H of $k^{n \times p}$ whose maximum rank is $< \min(n, p)$ is called a *subspace of matrices of low rank*. Such a subspace always comes from some linear matrix (for the M_i , take a spanning set of H).

These subspaces have been considered by several authors [F], [W], [AL], [A], [B], [EH], [Re]. They are not completely well-understood. Among them, the simplest one are the so called *compression space* (the terminology is from Eisenbud and Harris; Westwick [W] calls *essentially decomposable* such a space, in the case where all nonzero matrices in H have the same rank r): H is a compression space if after some change of basis of rows and columns over k (independently), the matrix M has the block form $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$, where the matrix B is of size $i \times j$, and where moreover the maximum rank in H is $i + j$ (note that a matrix of this form, with B of size $i \times j$, has necessarily rank $\leq i + j$).

Our characterization is the following (Theorem 3.1): *H is a compression space if and only if the commutative and noncommutative rank of M are equal.* Before proving this result, we need to generalize a result of Cohn that characterizes linear square matrices that are invertible in the free field. We characterize the noncommutative rank of any linear matrix (see Theorem 2.1 for a precise statement). As a corollary, we obtain that the noncommutative rank of a linear matrix $M = M_0 + \sum_{i=1}^d x_i M_i$ depends only on the subspace H spanned by the M_i .

In the last section, we give an efficient algorithm to solve the following question: given a subspace H of $k^{n \times p}$ of low rank, decide if H is a compression space (note that in order to know that H is of low rank, it is enough to compute the commutative rank of M , which is easy), see Theorem 4.1. Note that in [EH] is given an effective criterion for a subspace H to be a compression space, but the underlying algorithm seems to be of high complexity. Our algorithm uses only techniques of linear algebra.

From Theorem 3.1 and Theorem 4.1 we may deduce two proofs of the following fact: the property for H to be a compression space or not is invariant under extension of the field of scalars k . This is not so obvious because, translating the definition into equations leads to a system of nonlinear algebraic equations.

2. RANK OVER THE FREE FIELD OF LINEAR MATRICES

Let X be a set of noncommuting variables and k a (commutative) field. We denote by $k\langle X \rangle$ the k -algebra of noncommutative polynomials over k generated by the noncommuting variables $x \in X$. Among all the fields containing $k\langle X \rangle$, there is one, called the *free field*, which is unique up to isomorphism and which is characterized by the following property: each square matrix M over $k\langle X \rangle$, which is full, becomes invertible over the free field. Recall that a square matrix M over a ring R is called *full* (over R) if it is not possible to have a factorization $M = PQ$, with P of size $n \times p$, Q of size $p \times n$ and $p < n$. Observe that the embedding of $k\langle X \rangle$ in the free field inverts the maximum possible matrices over $k\langle X \rangle$, since a non full matrix cannot be invertible, in any extension field of $k\langle X \rangle$.

More generally, define the *inner rank* of an $n \times p$ matrix M over a ring R to be the least r such that M has a factorization $M = PQ$, with P of size $n \times r$ and Q of size $r \times p$. Then a fundamental result of Cohn is the following: the inner rank of any matrix over $k\langle X \rangle$ is equal to its

rank over the free field (see [C1] p. 249–250). We denote the free field by $k\langle X \rangle$. See [C1], [C2] for more on the free field.

We consider now linear matrices, that is, matrices whose entries are polynomials of degree ≤ 1 . These matrices play a special role for the free field, since each element of the free field is equal to some entry of the inverse in the free field of some square linear matrix [C2], Theorem 6.3.7.

Recall from the Introduction that to each linear matrix over $k\langle X \rangle^{n \times p}$, we associate a subspace of $k^{n \times p}$.

We say that two subspaces H, K of $k^{n \times p}$ are *equivalent* if $K = UHV$ for some invertible matrices U, V over k ; equivalently, H is obtained from K by row and column operations over k . Now, Atkinson and Lloyd [AL] call *r-decomposable* a subspace H of $k^{n \times p}$ if H is equivalent to subspace of matrices all of the form $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$, where B is of size $i \times j$ and $i + j = r$ (equivalently, the 0 block is of size $n - i \times p - j$). Note that such a matrix is necessarily of rank $\leq r$. Following them, we say that a linear matrix M is *r-decomposable* if its associated subspace is; equivalently, for some invertible matrices U, V over k , UMV is of the above form.

The next result characterizes the rank of a linear matrix by a linear – algebraic property. It extends Cohn’s characterization of full linear matrices [C2] Cor. 6.3.6.

Theorem 1. *Let M be a linear matrix of size $n \times p$ over $k\langle X \rangle$ and $r < \min(n, p)$. Its rank in the free field (equivalently, its inner rank in $k\langle X \rangle$) is $\leq r$ if and only if M is *r-decomposable*.*

Note that if the rank of a linear matrix is not $< \min(n, p)$ then it is equal to $\min(n, p)$: thus the theorem completely characterizes the rank of a linear matrix in the free field.

Proof. 1) Note that

$$\begin{pmatrix} A & 0 \\ B & C \end{pmatrix} = \begin{pmatrix} A & 0 \\ B & I_i \end{pmatrix} \begin{pmatrix} I_j & 0 \\ 0 & C \end{pmatrix}$$

where I_l denotes the identity matrix of size $l \times l$; the product is well-defined, since A has j columns, and C has i rows. In particular, the number of columns of the first factor (= number of rows of the second) is $j + i = r$, hence the product has inner rank $\leq r$.

2) In order to prove the converse, we may suppose that the rank of M in the free field is exactly r . Then we may write $M = FG$, where

F, G are matrices over $k\langle X \rangle$ of size $n \times r$ and $r \times p$. Since $r < n$, the rank in the free field of F is $\leq r$; it is actually exactly r , otherwise M has rank $< r$ in the free field, a contradiction. This implies that F is right regular, i.e. $FH = 0 \Rightarrow H = 0$ (otherwise, F has a kernel and its rank is $< r$). Symmetrically, G is left regular. Since F is right regular and G is left regular, and since $M = FG$ is of degree ≤ 1 we may apply Lemma 6.3.4 of [C2]: hence we may assume that $\deg(F) \leq 1$.

Suppose that $\deg F = 0$. This implies that the rows of M are k -linear combinations of the r rows of G ; thus the rows of M are of rank $\leq r$ over k , and by row operations over k , we may annihilate $n - r$ of them: we obtain a rectangle of 0's of size $(n - r) \times p$, which proves the theorem in this case.

We assume now that $\deg F = 1$. Then by [C2], Corollary VI.3.3, there exist $U \in GL_n(k)$ and $W \in GL_r(k\langle X \rangle)$ such that

$$UFW = \begin{pmatrix} A & 0 \\ 0 & I_s \end{pmatrix},$$

where A is *left monic*: this means that $A = A_0 + \sum_{x \in X} xA_x$, $A_0, A_x \in k^{n-s \times (r-s)}$, and the rows of the $A_x, x \in X$, span k^{r-s} .

Observe that this latter condition implies that

$$(2.1) \quad \deg(AN) = 1 + \deg(N)$$

for each matrix N over $k\langle X \rangle$; indeed, we may assume that N is homogeneous of degree d , and write $N = \sum wN_w$, where the sum is over all words w on X of length d and where the N_w are matrices over k . Then the term of degree $d + 1$ in AN is $\sum_{x,w} xwA_xN_w$. If it is zero, then $A_xN_w = 0$, and since the rows of the A_x span k^{r-s} , $N_w = 0$, a contradiction since N is of degree d . Hence the term of degree $d + 1$ in AN is nonzero, and this proves (1).

We have $UM = UFG = UFWW^{-1}G$. Let us partition $W^{-1}G = \begin{pmatrix} G_1 \\ G_2 \end{pmatrix}$, so that G_1, G_2 are of size $(r - s) \times p, s \times p$. Then

$$UM = \begin{pmatrix} A & 0 \\ 0 & I_s \end{pmatrix} \begin{pmatrix} G_1 \\ G_2 \end{pmatrix} = \begin{pmatrix} AG_1 \\ G_2 \end{pmatrix}.$$

This implies that AG_1, G_2 are linear since UM is. Moreover, by (1), G_1 must be of degree 0. Since G is left regular and W invertible, $W^{-1}G$ is also left regular, and this implies that G_1 also is. Thus the rank over k of $G_1 \in k^{(r-s) \times p}$ is $r - s$ (in particular $r - s \leq p$), and by column operations over k we may bring G_1 to the form $(I_{r-s}, 0_{r-s, p-r+s})$; in other words,

there exists $V \in GL_p(k)$ such that $G_1V = (I_{r-s}, 0_{r-s, p-r+s})$. We obtain finally

$$W^{-1}GV = \begin{pmatrix} G_1 \\ G_2 \end{pmatrix} V = \begin{pmatrix} I_{r-s} & 0_{r-s, p-r+s} \\ B & C \end{pmatrix}$$

and, since $UMV = UFWW^{-1}GV$,

$$UMV = \begin{pmatrix} A & 0 \\ 0 & I_s \end{pmatrix} \begin{pmatrix} I_{r-s} & 0_{r-s, p-r+s} \\ B & C \end{pmatrix} = \begin{pmatrix} A & 0_{n-s, p-r+s} \\ B & C \end{pmatrix}$$

since A has $n - s$ rows; this concludes the proof. \square

Corollary 1. *The rank of a linear matrix $M = M_0 + \sum_{i=1}^d x_i M_i$ over the free field $k \langle x_1, \dots, x_d \rangle$ depends only on the subspace of $k^{n \times p}$ spanned by the matrices M_i .*

Hence this rank gives an invariant of subspaces of matrices. It seems not easy to calculate. The algorithm of [CR] (which decides if a linear matrix is full) may be easily adapted to compute the rank of a linear matrix over the free field; it is however of high complexity, since it uses Grbner bases. It would be interesting to find an algorithm which uses only linear algebra techniques.

The following result compares the commutative and noncommutative ranks.

Corollary 2. *Let M be a linear matrix. Then $crk(M) \leq ncrk(M) \leq 2crk(M)$.*

Proof. The first inequality follows from the theorem.

For the second, we use a result of [F] (Lemma 1): if a subspace H of $k^{n \times p}$ has maximum rank r , then it is equivalent to a subspace, all matrices of which are of the form $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$, where B is square of order r . Hence, we may suppose that M is of this form. Thus the theorem implies that $ncrk(M) \leq 2r$. \square

Remark 1. The first inequality in the corollary is evidently sharp: indeed, take a linear matrix of size $n \times p$ and of commutative rank $\min(n, p)$. However, the second is not. Indeed, if M has commutative rank 1, then necessarily it has also noncommutative rank 1; this is because, classically, such a matrix is equivalent (after change of bases

over k in the spaces of rows and of columns) to a matrix of the form

$$\begin{pmatrix} 0 & \cdots & 0 & * \\ 0 & \cdots & 0 & * \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & * \end{pmatrix}$$

or its transpose.

The case of rank 2 and 3 may also be handled by results of Atkinson [A] and Eisenbud-Harris [EH]. Suppose that M is of commutative rank 2 and let H be the span in $k^{n \times p}$ of the M'_i 's. Then it follows from Theorem 1.1 in [EH] that its noncommutative rank is 2 or 3 (since the matrix in the Introduction has noncommutative rank 3).

Similarly, suppose that M has commutative rank 3. Then it follows from Theorem 1.2 in [EH] that M has noncommutative rank 3 or 4.

Note that, by direct sum of the matrix of the Introduction, one may find a square linear matrix of order $3n$, which is of noncommutative rank $3n$, and of commutative rank $2n$. So, one could expect an inequality of the form $ncrk(M) \leq \frac{3}{2}crk(M)$, which would certainly be sharp.

3. COMPRESSION SPACES

We recall the definitions of the Introduction, and we choose the language of linear mappings, instead of matrices.

We consider a subspace H of $\text{Hom}(E, F)$, where E, F are vector spaces over k , of dimension n, p respectively. We assume that k is infinite.

By definition, the *rank* of H is the maximum rank of its elements. We say that H is of *low rank* if its rank is smaller than $\min(n, p)$.

Among subspaces of low rank are the following ones: H is a *compression space* if for some subspaces E' of E and F' of F one has:

- 1) $\text{codim } E' + \dim F' = \text{rank of } H$;
- 2) any element of H maps E' into F' .

Taking bases of E and F , containing bases of E' and F' , we see that the matrices representing H are of all the form $\begin{pmatrix} \times & 0 \\ \times & \times \end{pmatrix}$, where the 0 matrix has $\text{codim } F'$ lines and $\dim E'$ columns. In other words, in the terminology of [AL], H is a compression space if H is r -decomposable and of rank r .

Lemma 1. *Let $M = M_0 + \sum_{i=1}^d x_i M_i$ be a linear matrix, and H the subspace spanned by the matrices M_0, M_1, \dots, M_d . Then the maximum rank in H is equal to the commutative rank of M (k is infinite).*

Proof. Note that, for square matrices A_i over k , $\det \left(x_0 A_0 + \sum_{i=1}^d x_i A_i \right)$ vanishes if and only so does $\det \left(A_0 + \sum_{i=1}^d x_i A_i \right)$, since the first is an homogeneous polynomial of degree the order of the A_i 's. If we apply this to the minors of M , we see that it is enough to prove the lemma for the linear matrix $\sum_{i=0}^d x_i M_i$. Now, since k is infinite, a polynomial vanishes if and only if it vanishes for all values of the variables in k . This implies the lemma. \square

Given a subspace H of $\text{Hom}(E, F)$, viewed in matrix form, we associate to H a linear matrix as follows: let M_1, \dots, M_d be a spanning set of H , let x_1, \dots, x_d be indeterminates; then the linear matrix is
$$M = \sum_{i=1}^d x_i M_i.$$

We know from Lemma 3.1 that $\text{crk}(M) = \text{rank}(H)$. Suppose that H is not of low rank; equivalently, its rank is $\min(n, p)$. Then, since a $n \times p$ matrix over any field has $\text{rank} \leq \min(n, p)$, we deduce that $\text{crk}(M) = n \text{crk}(M)$ if H is not of low rank.

The striking fact is that, when H is of low rank, then this equality characterizes compression spaces.

Theorem 2. *Let H be a space of matrices of low rank and M its associated linear matrix. Then H is a compression space if and only if the commutative rank of M and its noncommutative rank coincide.*

Proof. Suppose that $\text{crk}(M) = r = n \text{crk}(M)$. Note that $r < \min(n, p)$ since H is of low rank. Then by Theorem 2.1, after suitable change of bases in E, F we may assume that M is of the form $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$, where the zero matrix is of size $(n - i) \times (p - j)$, with $i + j = r$. Let E' be the subspace of E spanned by the last $p - j$ basis vectors, and F' the subspace of F spanned by the last i basis vectors. Then we see that each element of H maps E' into F' . Moreover, $\text{codim } E' + \dim F' = j + i = r$. Hence H is a compression space.

Conversely, if H is a compression space, let r be its rank. Then, by definition, we may after change of basis in E, F bring H into the form $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$, where A has $\text{codim } E'$ columns, and C has $\dim F'$ rows, with $r = \text{codim } E' + \dim F'$. Then M is also of this form, and its

noncommutative rank is $\leq r$ by Theorem 2.1. Hence, by Corollary 2.2, it is exactly r . \square

Corollary 3. *Let $k \subset k'$ be an extension of commutative fields. Let H be a vector space of matrices of low rank over k , and H' its extension to k' . Then H is a compression space if and only if H' is a compression space.*

Proof. Let M be the associated linear matrix; M has coefficients in k . Its commutative rank is unchanged under the extension $k \subset k'$. Furthermore, its noncommutative rank is also unchanged, since the free field $k \langle x_1, \dots, x_d \rangle$ embeds in the free field $k' \langle x_1, \dots, x_d \rangle$ (see [C2] Theorem 6.4.6). Thus, the corollary follows from the theorem. \square

A more elementary proof of this corollary will be given in Section 4.

4. AN ALGORITHM

Let E, F be vector spaces of dimension n, p and let $H \subset \text{Hom}(E, F)$ be a subspace of low rank $r < \min(n, p)$. Select an $f \in H$ with $\text{rank}(f) = r$.

Define the sequence of subspaces (E_i) of E and (F_i) of F by: $F_0 = \{0\}$, and for $i \geq 1$, recursively,

$$E_i = f^{-1}(F_{i-1}), \quad F_i = \sum_{g \in H} g(E_i). \quad (*)$$

For effective computations, note that, if H is given, by a basis (g_j) for instance, one has $F_i = \sum_j g_j(E_i)$, so that the sequence may effectively be computed.

Note that both sequences $(F_i), (E_i)$ are increasing, since $F_0 = \{0\} \subseteq F_1$, and: $F_{i-1} \subseteq F_i \Rightarrow E_i \subseteq E_{i+1}$ and $F_i \subseteq F_{i+1}$.

Thus, there is some p such that $F_{p-1} = F_p$, and for this p one has: $E_p = f^{-1}(F_{p-1}) = f^{-1}(F_p)$, thus $f^{-1}(F_p) = E_p$ and $\forall g \in H, g(E_p) \subseteq F_p$.

Theorem 3. *With the previous notations, H is a compression space if and only if $\text{codim } E_p + \dim F_p = r$.*

Proof. If this last equality holds, surely H is a compression space, since each g in H maps E_p into F_p .

Conversely, suppose that H is a compression space. Then for some subspace E', F' of E, F , we have $\text{codim } E' + \dim F' = r$, and each element of H maps E' into F' . In particular $f(E') \subseteq F'$.

We claim that $f(E') = F'$ and $f^{-1}(F') = E'$. Indeed let \bar{f} be the composition $E \xrightarrow{f} F \longrightarrow F/F'$, where the second mapping is the canonical one. Then $\text{Ker } \bar{f} = f^{-1}(F')$. We have $E' \subseteq \text{Ker } \bar{f}$, since $f(E') \subseteq F'$, so that $\dim E' \leq \dim(\text{Ker } \bar{f})$.

Now, $\text{rank}(f) = r$, hence $\text{rank}(\bar{f}) \geq r - \dim F'$ with equality if and only if $F' \subseteq f(E)$. We have by hypothesis $r - \dim F' = \text{codim } E' = \dim E - \dim E' = \text{rank}(\bar{f}) + \dim(\text{Ker } \bar{f}) - \dim E'$; hence the previous inequality implies $\dim(\text{Ker } \bar{f}) \leq \dim E'$, which implies equality and $E' = \text{Ker } \bar{f} = f^{-1}(F')$. Using $\dim(\text{Ker } \bar{f}) = \dim E'$, the same computation shows that $\text{rank}(\bar{f}) = r - \dim F'$. This implies by a previous remark that $F' \subseteq f(E')$ and finally, that $f(E') = F'$.

Observe that $F_0 \subseteq F'$. If $F_{i-1} \subseteq F'$, then by the claim $E_i = f^{-1}(F_{i-1}) \subseteq f^{-1}(F') = E'$, and $F_i = \sum_{g \in H} g(E_i) \subseteq \sum_{g \in H} g(E') \subseteq F'$.

Thus, we obtain by induction that $E_i \subseteq E'$, and $F_i \subseteq F'$ for each i .

This is true in particular for $i = p$, so that, changing notations ($A = E_p, B = F_p$), we have a subspace A of E' and a subspace B of F' such that: $\forall g \in H, g(A) \subseteq B$ and $f^{-1}(B) = A$.

We show that $\text{codim } A + \dim B = r$, which will prove the theorem. Choose a subspace A' of E' such that $A \oplus A' = E'$. We claim that the composition $u : A' \xrightarrow{f} F' \longrightarrow F'/B$ is an isomorphism.

Taking the claim for granted, we obtain $\dim A' = \dim F' - \dim B$, thus $r = \text{codim } E' + \dim F' = \dim E - \dim E' + \dim F' = \dim E - \dim A - \dim A' + \dim F' = \dim E - \dim A + \dim B = \text{codim } A + \dim B$.

Let us prove the claim. We have $f(A) \subseteq B$. In fact, $f(A) = B$; indeed, if $b \in B$, then, since $f(E') = F', b = f(e')$ for some $e' \in E'$; then $e' \in f^{-1}(b) \subseteq f^{-1}(B) = A$, hence $e' \in A$ and $b \in f(A)$, which shows that $B \subseteq f(A)$.

Thus we have $f(A) = B \Rightarrow F' = f(E') = f(A) + f(A') = B + f(A')$ and this implies that the restriction to $f(A')$ of the canonical mapping $F' \rightarrow F'/B$ is surjective; in other words, u is surjective.

Now, u is also injective, since $\text{Ker } u = f^{-1}(B) \cap A' = A \cap A' = \{0\}$. \square

Now, the algorithm works as follows. By using the linear matrix M associated to H , compute the maximal rank r of H ; by finding in the infinite field k values of the variables appearing in M that do not annihilate some nonzero $r \times r$ minor of M , one obtains an element f in H of rank r . Then one computes the subspaces E_i, F_i described at the beginning of the section; note that this computation is of low complexity, since it amounts to solve linear equations. Then one finds

$p \leq \dim F$ such that $F_{p-1} = F_p$ as above. It is enough then to check if $\text{codim } E_p + \dim F_p = r$ and apply the theorem.

Note that in [EH], an effective criterion is given for a space H of matrices of low rank to be a compression space; however, the computations use Gröbner bases, so are of high complexity (see [EH] p. 150–151).

Second proof of Corollary 3.1. The spaces E_i, F_i are all defined over k , since so are H and f . Moreover, $F_{p-1} = F_p$ if and only if $F_{p-1} \otimes_k k' = F_p \otimes_k k'$, and the condition of the theorem is invariant under extension, since the rank of H does not change under commutative field extension. This proves the corollary. \square

REFERENCES

- [A] M.D. Atkinson, Primitive spaces of matrices of bounded rank II, *J. Austral. Math. Soc. Ser. A* 34, 1983, 306–315.
- [AL] M.D. Atkinson, S. Lloyd, Large spaces of matrices of bounded rank, *Quarterly Journal of Mathematics Oxford* 31, 1980, 253–262.
- [B] L.B. Beasley, *Null spaces of matrices of bounded rank*, Current trends in matrix theory, 45–50, North Holland, New York, 1987.
- [C1] P.M. Cohn, *Free rings and their relations*, Academic Press, 1985 (first edition 1971).
- [CR] P.M. Cohn, C. Reutenauer, On the construction of the free field, *Intern. J. Alg. Computation* 9, 1999, 307–323.
- [C2] P.M. Cohn, *Skew fields, theory of general division rings*, Cambridge University Press, 1995.
- [C3] P.M. Cohn, The word problem for free fields, *J. Symb. Logic* 38, 1973, 309–314; correction and addendum 40, 1975, 69–74.
- [E] S. Eilenberg, *Automata, languages and machines*, vol. A, Academic Press, 1974.
- [EH] D. Eisenbud, J. Harris, Vector spaces of matrices of low rank, *Advances Math.* 70, 1988, 135–155.
- [F] H. Flanders, On spaces of linear transformations with bounded rank, *J. London Math. Soc.* 37, 1962, 10–16.
- [G] F.R. Gantmacher, *Applications of the theory of matrices*, Interscience, 1959.
- [H] J. Harris, *Algebraic geometry, a first course*, Springer, 1993.
- [K] D.E. Knuth, *The art of computer programming*, vol. 2, Seminumerical problems, Addison Wesley, 1981.
- [Re] R. Re, A note on linear subspaces of determinantal varieties, *Le Matematiche* 50, 1995, 173–178.
- [Ro] M.L. Roberts, Endomorphism rings of finitely presented modules over free algebras, *J. London Math. Soc.* 30, 1984, 197–209.
- [W] R. Westwick, *Spaces of linear transformations of equal rank*, Linear Algebra and Its Applications 5, 1972, 49–64.