

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

European Journal of Combinatorics

journal homepage: www.elsevier.com/locate/ejc

Primitive elements of the Hopf algebras of tableaux

C. Malvenuto^a, C. Reutenauer^b^a Dipartimento di Matematica "Guido Castelnuovo", Sapienza Università di Roma, Italy^b Département de mathématiques, Université du Québec à Montréal, Canada

ARTICLE INFO

Article history:

Received 27 November 2020

Accepted 17 May 2021

Available online xxxx

ABSTRACT

The character theory of symmetric groups, and the theory of symmetric functions, both make use of the combinatorics of Young tableaux, such as the Robinson–Schensted algorithm, Schützenberger’s “*jeu de taquin*”, and evacuation. In 1995 Poirier and the second author introduced some algebraic structures, different from the plactic monoid, which induce some products and coproducts of tableaux, with homomorphisms. Their starting point are the two dual Hopf algebras of permutations, introduced by the authors in 1995. In 2006 Aguiar and Sottile studied in more detail the Hopf algebra of permutations: among other things, they introduce a new basis, by Möbius inversion in the poset of weak order, that allows them to describe the primitive elements of the Hopf algebra of permutations. In the present Note, by a similar method, we determine the primitive elements of the Poirier–Reutenauer algebra of tableaux, using a partial order on tableaux defined by Taskin.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

In 1995 the authors of the present paper introduced two dual Hopf algebra structures on permutations [11]. The products and coproducts of permutations originated from the concatenation Hopf algebra and shuffle Hopf algebra on $\mathbb{Z}\langle\mathbb{N}^{>0}\rangle$, the module generated by words of positive integers, from Solomon’s descent algebra [15] and Gessel’s (internal) coalgebra [5] of quasi-symmetric functions. The two Hopf structures on $\mathbb{Z}S$, the module with \mathbb{Z} -basis all the permutations

E-mail addresses: claudia@mat.uniroma1.it (C. Malvenuto), Reutenauer.Christophe@uqam.ca (C. Reutenauer).

in $S = \cup_{n \geq 0} S_n$, for S_n the symmetric group on $\{1, \dots, n\}$, are self-dual (this result is somewhat implicit in the earlier work of the authors of the present article, but is explicit in [7] and [2]).

In the same year [13], carrying on these themes, Poirier and the second author proved that the two dual Hopf structures on $\mathbb{Z}S$ are free associative algebras. By restriction on the plactic classes they obtained two dual structures of Hopf algebras on the \mathbb{Z} -module $\mathbb{Z}S$ with basis the set T of all standard Young tableaux. The product and coproduct are described there in terms of Schützenberger’s “*jeu de taquin*” [9]. They also provided different morphisms between these structures and the descent algebras, symmetric functions and quasi-symmetric functions. In particular, the map sending a permutation into its left tableau in the Schensted correspondence is a Hopf morphism.

Loday and Ronco [10] characterized the product of two permutations by the use of the weak order of permutations: it is the sum of all permutations in some interval for this order. In 2005 [1], Aguiar and Sottile studied in further and thorough details the structure of the Hopf algebras of permutations, giving explicit formulas for its antipode, proving that it is a cofree algebra and determining its primitive elements. For the latter task, they introduced a new basis of $\mathbb{Z}S$, related to the basis of permutations via Möbius inversion in the poset of the Bruhat weak order of the symmetric groups. In [4], Duchamp, Hivert, Novelli and Thibon studied the Hopf algebra of permutations (denoted there $FQSYM$), and gave among others a faithful representation by noncommutative polynomials.

The Hopf algebra of tableaux was used by Jöllenbeck [7], and Blessenohl and Schocker [2], to define their noncommutative character theory of the symmetric group. Moreover, Muge Taşkın [16] used the order on tableaux, induced by the weak order of permutations, to characterize the product of two tableaux, in a way reminiscent of the result of Loday and Ronco.

The purpose of this Note is to find the primitive elements of $\mathbb{Z}T$, the Hopf algebra of tableaux with respect to the product and coproduct, following the approach of Aguiar and Sottile. A new basis for $\mathbb{Z}T$ is obtained by Möbius inversion for the Taşkın order of tableaux. The nice feature of the proofs here is that we manage to avoid “*jeu de taquin*”, and use a simpler description through a shifted left and right concatenation product of tableaux.

Note that Taşkın studies in [16] four orders on tableaux, which appeared in earlier work of Melnikov [12]; she calls the weak order on tableaux the *induced Duflo order*. The four orders are mutually comparable, and the weak order is the smallest one. Moreover, Taşkın, with Karaali and Senturia, introduced recently a new order on Young tableaux [8]. We did not explore what happens if one performs an change of basis based on these orders (except the weak order). It could be that the primitive elements of $\mathbb{Z}T$ behave nicely in this new basis, and that moreover the multiplicative constants in one of these basis are positive; but this is open.

2. Preliminaries on permutations

We denote by S_n symmetric group on $\{1, \dots, n\}$. We often represent permutations as words: $\sigma \in S_n$ is represented as the word $\sigma(1)\sigma(2) \dots \sigma(n)$. By abuse of notation, we identify σ and the corresponding word. A *word* in the sequel will always be on the alphabet of positive integers, also called *letters*. We denote by $|\sigma|$ the number of letters of σ .

We denote by \leq the *right weak order* of permutations. Recall that it is defined as the reflexive and transitive closure of the relation $u < v$, $u, v \in S_n$, $v = u \circ \tau$, for some adjacent transposition $\tau \in S_n$ such that $l(u) < l(v)$, where $l(u)$ denotes as usual the *length* of u in the sense of Coxeter groups. Recall that this order may also be defined by comparing inversions sets: let $Inv(\sigma)$ be the set of inversions (‘by values’) of σ , that is, the set of pairs (j, i) with $j > i$ and $\sigma^{-1}(j) < \sigma^{-1}(i)$; then $u \leq v$ if and only if $Inv(u) \subseteq Inv(v)$. Note that (j, i) is an inversion of σ if and only if $j > i$ and j appears on the left of i in the word representing σ . This definition applies also to any word.

Given $\sigma \in S_n$, and a subset I of $[n]$, $\sigma \setminus I$ denotes the word obtained by removing in the word σ the digits not in I (whereas the restriction of σ to I is the subword $\sigma|_I$ of the images of the letters in I). For example, for $\sigma = 2517643$, one has $2517643 \setminus \{2, 3, 6\} = 263$ (while $\sigma|_{\{2,3,6\}} = 514$).

Moreover, v being a word without repetition of letters, $st(v)$ denotes the *standard permutation* of the word v , obtained by replacing each letter in v by its image under the unique increasing bijection from the set of letters in v onto $\{1, 2, \dots, |v|\}$. For example, $st(5713) = 3412$.

Lemma 2.1. *If $s \leq t$ are permutations in S_n and I an interval in $\{1, \dots, n\}$, then $st(s | I) \leq st(t | I)$.*

Proof. Let (j, i) be an inversion of $s | I$; then $i, j \in I$. Then (j, i) is an inversion of s , hence of t . It follows that it is also an inversion of $t | I$. Since standardizing amounts to apply an increasing bijection, we deduce the lemma. \square

Let $S = \bigcup_{n \geq 0} S_n$ be the disjoint union of all symmetric groups. There is a classical associative product on S , denoted by \square , which turns it into a free monoid [13]: let $u \in S_p$ and $v \in S_q$; let \bar{v} be obtained by adding p to each digit in v ; then $u \square v$ is the concatenation $u\bar{v}$ of u and \bar{v} (or *right shifted concatenation*). For example, $231 \square 12 = 23145$, with here $p = 3$. The free generators of this free monoid are the *indecomposable* permutations, which have some importance in algebraic combinatorics; see [3].

A variant of this product is as follows: given two permutations as above, $v \Delta u = \bar{v}u$ (which we refer to as *left shifted concatenation*). Example: $12 \Delta 231 = 45231$.

Clearly, the product \square and the opposite product of Δ are conjugate under the mapping $w \mapsto \tilde{w}$ which reverses words:

$$v \Delta u = \widetilde{u \square v}.$$

It follows that S with the product Δ is also a free monoid, freely generated by the permutations which are indecomposable for this product, which are the reversals of the indecomposable permutations.

For later use, we need

Lemma 2.2. *The weak order \leq on permutations is compatible with the product Δ : $u \leq u', v \leq v' \Rightarrow v \Delta u \leq v' \Delta u'$.*

Proof. Let $u \in S_p, v \in S_q$. Suppose that $u \leq u', v \leq v'$. We show that $v \Delta u \leq v' \Delta u'$. Let (j, i) be an inversion of $v \Delta u = \bar{v}u$. If $i, j \leq p$, then by construction of the product Δ , (j, i) is an inversion of u , hence of u' (since $u \leq u'$), hence of $v' \Delta u'$. If $i, j > p$, then $(j - p, i - p)$ is an inversion of v , hence of v' (since $v \leq v'$) and therefore (j, i) is an inversion of $v' \Delta u'$. If $i \leq p$ and $j > p$, then (j, i) is an inversion of $v' \Delta u'$. There is no other case, since $i < j$. \square

3. Hopf algebra of permutations

Denote by $\mathbb{Z}S$ be the free \mathbb{Z} -module with basis S . We define on $\mathbb{Z}S$ a product, denoted by $*$ (called *destandardized concatenation*), and a coproduct (called *standardized unshuffling*), denoted by δ , which turn it into a Hopf algebra (see [11]). If $\alpha \in S_p, \beta \in S_q, \alpha * \beta$ is the sum of all permutations in S_{p+q} of the form uv (concatenation of u and v), where u, v are of respective lengths p, q and $st(u) = \alpha, st(v) = \beta$; for example, $12 * 21 = 1243 + 1342 + 1432 + 2341 + 2431 + 3421$. Moreover, for $\sigma \in S_n$,

$$\delta(\sigma) = \sum_{0 \leq i \leq n} \sigma | \{1, \dots, i\} \otimes st(\sigma | \{i + 1, \dots, n\}).$$

An example of coproduct is $\delta(3124) = \epsilon \otimes st(3124) + 1 \otimes st(324) + 12 \otimes st(34) + 312 \otimes st(4) + 3124 \otimes \epsilon = \epsilon \otimes 3124 + 1 \otimes 213 + 12 \otimes 12 + 312 \otimes 1 + 3124 \otimes \epsilon$. Here ϵ is the empty permutation in S_0 , the neutral element of the bialgebra $\mathbb{Z}S$.

Following Aguiar and Sottile [1], we define a new linear basis \mathcal{M}_σ of $\mathbb{Z}S$, indexed by permutations, and called *monomial basis*. (Notice that Aguiar and Sottile deal with the isomorphic dual Hopf algebra, the isomorphism being $\sigma \mapsto \sigma^{-1}$, and also that they use the left weak order.) These elements are given by the formula

$$\sigma = \sum_{\sigma \leq w} \mathcal{M}_w,$$

which defines them recursively, via Möbius inversion formula, since \leq is an order.

The following result is equivalent to a result due to Aguiar and Sottile ([1] Theorem 3.1).

Theorem 3.1. For any permutation σ , one has

$$\delta(\mathcal{M}_\sigma) = \sum_{\sigma=v\Delta u} \mathcal{M}_u \otimes \mathcal{M}_v.$$

We give below the proof of this result; it may help to understand the proof of the similar result on the Hopf algebra of tableaux, which we give in Section 6.

We begin by a lemma, that will also have an analogue for tableaux.

Lemma 3.1. Let $n = p + q$, $\sigma \in S_n$, $a = \sigma | \{1, \dots, p\}$, $b = st(\sigma | \{p + 1, \dots, n\})$. Then, for $v \in S_q$, $u \in S_p$, $\sigma \leq v \Delta u$ if and only if $a \leq u$ and $b \leq v$.

Proof. 1. We show first that $\sigma \leq b \Delta a$. We have $b \Delta a = \bar{b}a$ (concatenation of words), where it is easily seen that $\bar{b} = \sigma | \{p + 1, \dots, n\}$ (add p to each letter of b).

Let (j, i) be an inversion of σ . Then $i < j$ and j is at the left of i in σ . If $j, i \leq p$, then (j, i) is an inversion of a , and therefore also of $b \Delta a$. If $j, i > p$, then (j, i) is an inversion of \bar{b} , hence of $b \Delta a$. If $i \leq p$ and $j > p$, then (j, i) is an inversion of $b \Delta a$, since in this latter permutation, each letter $> p$ is at the left of each letter $\leq p$. There is no other case since $j > i$.

Thus (j, i) is in each case an inversion of $b \Delta a$, hence $\sigma \leq b \Delta a$.

2. Suppose that $a \leq u$ and $b \leq v$. Then clearly $b \Delta a \leq v \Delta u$, by Lemma 2.2. Hence by 1, $\sigma \leq v \Delta u$.

3. Suppose that $\sigma \leq v \Delta u$. If (j, i) is an inversion of a , then it is an inversion of σ , hence of $v \Delta u = \bar{v}u$; therefore it is an inversion of u since $i, j \leq p$; thus $a \leq u$. Similarly, $b \leq v$. \square

Proof of Theorem 3.1. Define the \mathbb{Z} -linear mapping $\delta_1 : \mathbb{Z}S \mapsto \mathbb{Z}S \otimes \mathbb{Z}S$ by $\delta_1(\mathcal{M}_\sigma) = \sum_{\sigma=v\Delta u} \mathcal{M}_u \otimes \mathcal{M}_v$. It is enough to show that $\delta_1 = \delta$. We have for any permutation $\sigma \in S_n$,

$$\delta_1(\sigma) = \delta_1\left(\sum_{\sigma \leq w} \mathcal{M}_w\right) = \sum_{\sigma \leq w} \sum_{w=v\Delta u} \mathcal{M}_u \otimes \mathcal{M}_v = \sum_{\sigma \leq v\Delta u} \mathcal{M}_u \otimes \mathcal{M}_v.$$

This is by Lemma 3.1 equal to

$$\begin{aligned} & \sum_{0 \leq i \leq n} \sum_{\substack{\sigma | \{1, \dots, i\} \leq u \\ st(\sigma | \{i+1, \dots, n\}) \leq v}} \mathcal{M}_u \otimes \mathcal{M}_v \\ &= \sum_{0 \leq i \leq n} \left(\sum_{\sigma | \{1, \dots, i\} \leq u} \mathcal{M}_u \right) \otimes \left(\sum_{st(\sigma | \{i+1, \dots, n\}) \leq v} \mathcal{M}_v \right) \\ &= \sum_{0 \leq i \leq n} \sigma | \{1, \dots, i\} \otimes st(\sigma | \{i + 1, \dots, n\}) = \delta(\sigma), \end{aligned}$$

by the definition of the basis \mathcal{M}_σ . \square

In order to understand the equivalence between the previous theorem and the theorem of Aguiar and Sottile, it may be useful to use the notion of global descents, introduced by them. Recall that according to Aguiar and Sottile ([1], Definition 2.12), a *global descent* of $\sigma \in S_n$ is a position $i \in \{1, \dots, n - 1\}$ such that for any $j \leq i$ and any $k > i$ one has $\sigma(j) > \sigma(k)$.

Then the permutations that are indecomposable for the product Δ (which are the free generators of the free monoid S) are those which have no global descents. Moreover, if $\sigma = \sigma_1 \Delta \dots \Delta \sigma_k$ with indecomposable σ_i 's, then the global descents of σ are the positions $|\sigma_1|, |\sigma_1| + |\sigma_2|, \dots, |\sigma_1| + \dots + |\sigma_{k-1}|$.

For example, $78\ 465\ 213 = 12 \Delta 132 \Delta 213$ has 2 and 5 as global descents, and 12, 132, 213 are indecomposable for Δ , equivalently, have no global descents.

Corollary 3.1. The submodule of primitive elements of $\mathbb{Z}S$ is spanned by the \mathcal{M}_σ such that σ has no global descent, or equivalently, σ is indecomposable for the product Δ .

4. Preliminaries on tableaux

For unreferenced results quoted here, see [14]. Denote by T_n denotes the set of standard Young tableaux (we say simply *tableaux*) whose entries are $1, \dots, n$. We denote by $(P(\sigma), Q(\sigma))$ the pair of tableaux associated with $\sigma \in S_n$ by the Schensted correspondence.

Let $T = \cup_{n \geq 0} T_n$ be the set of all standard tableaux. The *plactic equivalence* on T is the smallest equivalence relation generated by the *Knuth relations* $xjiky \sim_{plax} xjkij, xikjy \sim_{plax} xkijy, i < j < k$, for $i, j, k \in \mathbb{N}$ and $x, y \in \mathbb{N}^*$.

By Knuth's theorem, one has $P(\sigma) = P(\tau)$ if and only if $\sigma \sim_{plax} \tau$. Thus we may identify tableaux and plactic classes. In the sequel, we use systematically this identification.

Following Taskin [16], we define the *weak order* of tableaux as follows: let $U, V \in T_n$; let $u, v \in S_n$ be such that $P(u) = U, P(v) = V$. Define the relation $U \leq V$ if $u \leq v$; then \leq is the transitive closure of this relation.

In other words, the weak order on tableaux is the smallest order on T such that the mapping $P : S \rightarrow T$ is increasing for the weak order.

This order was introduced by Melnikov [12] (called there *Duflo order*) and Taskin [16] (denoted there \leq_{weak}). The difficulty here is to show that it is indeed an order.

For two tableaux A, U , one has $A \leq U$ if and only if there exist $n \geq 1$ and permutations $\alpha_0, \dots, \alpha_{n-1}, \beta_1, \dots, \beta_n$ such that

$$P(\alpha_0) = A, \alpha_0 \leq \beta_1 \sim \alpha_1 \leq \beta_2 \sim \dots \alpha_{n-1} \leq \beta_n, P(\beta_n) = U. \tag{1}$$

The product Δ of permutations induces a product of tableaux, still denoted by Δ . This follows from the next lemma.

Lemma 4.1. *The plactic equivalence is compatible with the product Δ , that is: $u \sim_{plax} u' \Rightarrow v \Delta u \sim_{plax} v \Delta u'$.*


Proof. Suppose indeed that $u, u' \in S_p, v \in S_q$ and for some letters $1 \leq i < j < k \leq p$, and words x, y , one has $u = xjiky, u' = xjkij$. Then $\bar{v}u = \bar{v}xjiky, \bar{v}u' = \bar{v}xjkij$ and therefore $v \Delta u \sim_{plax} v \Delta u'$. There are other similar cases, left to the reader. \square

Therefore, one has for any permutations u, v

$$P(v \Delta u) = P(v) \Delta P(u), \tag{2}$$

which means that P is a homomorphism from the monoid S into the monoid T , both with the product Δ .

Note that one may compute directly $V \Delta U$ as follows: $V \Delta U$ is the tableau obtained by letting fall \bar{V} onto U , with \bar{V} obtained by adding p to each letter in V (this follows from the dual Schensted correspondence, that is, column insertion). Thus $V \Delta U$ is the tableau denoted U/V in [16], p. 1109. For example:



We need also the following.

Lemma 4.2. *The weak order \leq on tableaux is compatible with the product Δ : $U \leq U', V \leq V' \Rightarrow V \Delta U \leq V' \Delta U'$.*

Proof. This follows from Lemma 2.2, and the characterization through Eq. (1) of the order, using the fact that P is an increasing surjective homomorphisms $S \rightarrow T$. \square

By Proposition 2.5 in [9], p.133, the plactic equivalence is compatible with the restriction to intervals, and with standardization. It follows that the plactic equivalence is compatible with the composition of the two operations: if $u, v \in S_n$, $u \sim_{plax} v$, and I is an interval of $[n]$, then $st(u | I) \sim_{plax} st(v | I)$. Thus, if $A = P(u)$, we may denote without ambiguity by $st(A | I)$ the tableau $P(st(u | I))$. By the work of Schützenberger, the corresponding tableau is obtained by jeu-de-taquin straightening of the skew tableau which is the restriction to I of the tableau $P(u)$; but we do not need this fact.

Lemma 4.3. *Let $A, B \in \mathcal{T}_n$ such that $A \leq B$, and I be an interval in $[n]$. Then $st(A | I) \leq st(B | I)$.*

Proof. This follows from the characterization through Eq. (1) of the order, Lemma 2.1, the previous observation, and the fact that P is increasing. \square

5. Main lemma

Recall the Taskin weak order on tableaux, denoted \leq .

Lemma 5.1. *Let $n = p + q$ and $\Sigma \in \mathcal{T}_n$. Let $A = st(\Sigma | \{1, \dots, p\})$, and let $B = st(\Sigma | \{p + 1, \dots, n\})$. Then for $U \in \mathcal{T}_p, V \in \mathcal{T}_q$, one has: $\Sigma \leq V \Delta U$ if and only if $A \leq U$ and $B \leq V$.*

Proof of Lemma 5.1. 1. We show first that $\Sigma \leq B \Delta A$. Let $\sigma \in S_n$ be such that $\Sigma = P(\sigma)$. Let $a = st(\sigma | \{1, \dots, p\})$ and $b = st(\sigma | \{p + 1, \dots, n\})$. Then $\sigma \leq b \Delta a$ by Lemma 3.1. It follows that $\Sigma \leq B \Delta A$ by Lemmas 4.1 and 4.2.

2. Suppose that $A \leq U$ and $B \leq V$. Then by 1. and Lemma 4.2, we have $\Sigma \leq B \Delta A \leq V \Delta U$.

3. Suppose now that $\Sigma \leq V \Delta U$. Let $u \in S_p, v \in S_q$ be such that $U = P(u), V = P(v)$. Then by Lemma 4.3, we have $A = st(\Sigma | \{1, \dots, p\}) \leq st((V \Delta U) | \{1, \dots, p\}) = P(st((v \Delta u) | \{1, \dots, p\})) = P(u) = U$. Moreover, by the same lemma,

$$\begin{aligned} B &= st(\Sigma | \{p + 1, \dots, n\}) \leq st((V \Delta U) | \{p + 1, \dots, n\}) \\ &= P(st((v \Delta u) | \{p + 1, \dots, n\})) = P(v) = V. \quad \square \end{aligned}$$

6. Primitive elements in the Hopf algebra of tableaux

The free \mathbb{Z} -module $\mathbb{Z}T$, based on the set T of tableaux, becomes a structure of Hopf algebra, quotient of the Hopf algebra $\mathbb{Z}S$ of Section 3, and whose product and coproduct are therefore also denoted by $*$ and δ . The quotient is obtained by identifying plactic equivalent permutations. In other words, consider the submodule I spanned by the elements $u - v, u \sim_{plax} v$; then I is an ideal and a co-ideal of $\mathbb{Z}S$, and the quotient $\mathbb{Z}S/I$ is canonically isomorphic with $\mathbb{Z}T$. Moreover, the canonical bialgebra homomorphism $\mathbb{Z}S \rightarrow \mathbb{Z}T$ maps each permutation σ onto $P(\sigma)$. See [13], Théorème 3.4 and 4.3 (iv), where the product and coproduct are there denoted $*'$ and δ' .

Now we introduce a new basis of $\mathbb{Z}T$, following the method of Aguiar and Sottile [1], replacing the weak order on permutations by the Taskin weak order on tableaux. The new basis (that we may call the *monomial basis*, following [1]) $\mathcal{M}_W, W \in T$, is completely defined by the identities

$$\Sigma = \sum_{\Sigma \leq W} \mathcal{M}_W,$$

for all tableau Σ , via Möbius inversion on the poset (T, \leq) .

Theorem 6.1. *Let $\Sigma \in \mathcal{T}_n$. Then*

$$\delta(\mathcal{M}_\Sigma) = \sum_{\Sigma = V \Delta U} \mathcal{M}_U \otimes \mathcal{M}_V.$$

Proof. Define the \mathbb{Z} -linear mapping $\delta_1 : \mathbb{Z}T \mapsto \mathbb{Z}T \otimes \mathbb{Z}T$ by

$$\delta_1(\mathcal{M}_W) = \sum_{W=V\Delta U} \mathcal{M}_U \otimes \mathcal{M}_V.$$

It is enough to show that $\delta_1 = \delta$.

We have

$$\begin{aligned} \delta_1(\Sigma) &= \delta_1\left(\sum_{\Sigma \leq W} \mathcal{M}_W\right) = \sum_{\Sigma \leq W} \sum_{W=V\Delta U} \mathcal{M}_U \otimes \mathcal{M}_V = \sum_{\Sigma \leq V\Delta U} \mathcal{M}_U \otimes \mathcal{M}_V \\ &= \sum_{p+q=n} \sum_{\substack{\Sigma \leq V\Delta U \\ V \in \mathcal{T}_q, U \in \mathcal{T}_p}} \mathcal{M}_U \otimes \mathcal{M}_V. \end{aligned}$$

This is by [Lemma 5.1](#), and with its notations, equal to

$$\begin{aligned} &\sum_{p+q=n} \left(\sum_{st(\Sigma | \{1, \dots, p\}) \leq U} \mathcal{M}_U \right) \otimes \left(\sum_{st(\Sigma | \{p+1, \dots, n\}) \leq V} \mathcal{M}_V \right) \\ &= \sum_{p+q=n} (st(\Sigma | \{1, \dots, p\})) \otimes (st(\Sigma | \{p+1, \dots, n\})). \end{aligned}$$

Choose σ such that $P(\sigma) = \Sigma$. Then by definition of $st(\Sigma | I)$, the latter quantity is equal to

$$\begin{aligned} &\sum_{p+q=n} P(st(\sigma | \{1, \dots, p\})) \otimes P(st(\sigma | \{p+1, \dots, n\})) \\ &= (P \otimes P)\left(\sum_{p+q=n} st(\sigma | \{1, \dots, p\}) \otimes st(\sigma | \{p+1, \dots, n\})\right) \\ &= (P \otimes P)(\delta(\sigma)) = \delta(P(\sigma)) = \delta(\Sigma), \end{aligned}$$

since P is a homomorphism of bialgebra, as recalled at the beginning of [Section 6](#). \square

Corollary 6.1. *The submodule of primitive elements of $\mathbb{Z}T$ is spanned by the \mathcal{M}_Σ such that Σ is indecomposable for the product Δ .*

The dimensions of the graded components of the submodule of primitive elements is therefore the sequence of the numbers of tableaux indecomposable for the product Δ ; it is denoted a_n in [\[13\]](#), p. 88–89. For $n = 1, 2, \dots, 10$, they are the numbers

$$1, 1, 1, 3, 7, 23, 71, 255, 911, 3535.$$

They appear as sequence A140456 in the Online Encyclopedia of Integer Sequences [\[6\]](#), with other interpretations.

7. Further remarks

7.1. Product formulas using Δ

The product Δ , both for permutations and tableaux, plays a role in product formulas in the dual Hopf algebras of $\mathbb{Z}S$ and $\mathbb{Z}T$.

First, one has to consider also the product \square of permutations: let $a \in S_p, b \in S_q$; then $a \square b = a\bar{b}$, where \bar{b} is obtained from b by adding p to each letter in b .

Recall the *shifted shuffle product* of permutations, denoted $\overline{\square}$: $a \overline{\square} b$ is the shuffle of a and \bar{b} . This product is the dual product of the coproduct δ of $\mathbb{Z}S$. On has

Theorem 7.1 (*Loday-Ronco [\[10\]](#) Theorem 4.1*). *Let $a, b \in S$. Then*

$$a \overline{\square} b = \sum_{a \square b \leq \sigma \leq b \Delta a} \sigma.$$


In other words, $a \square b$ is the sum of all permutations in the interval $[a \square b, b \triangle a]$ of the weak order.

The product \square is compatible with the plactic equivalence, hence \square is well-defined on tableaux. Denote also by \sqcup the product which is the dual of δ in the dual coalgebra of $\mathbb{Z}T$. Then one has

Theorem 7.2 (Taskin [16] Theorem 4.1). *Let $A, B \in \mathcal{T}$. Then*

$$A \square B = \sum_{A \square B \leq T \leq B \triangle A} T.$$

In other words, $A \square B$ is the sum of all tableaux belonging to the interval $[A \square B, B \triangle A]$ of the weak order. Note that the product $A \square B$ is denoted $A \sqcup B$ in [16], p. 1109. It corresponds to put the tableau \bar{B} aside the tableau A , then push every row of \bar{B} toward A . For example:



7.2. Multiplicative basis

Note that Theorem 6.1 means that in the dual Hopf algebra of $\mathbb{Z}T$, the dual basis \mathcal{M}_T^* of the basis \mathcal{M}_T is multiplicative, in the sense of [4]: one has for tableaux A, B ,

$$\mathcal{M}_A^* \sqcup \mathcal{M}_B^* = \mathcal{M}_{B \triangle A}^*.$$

Indeed, using the canonical pairing between $\mathbb{Z}T$ and its dual,

$$\begin{aligned} \langle \mathcal{M}_A^* \sqcup \mathcal{M}_B^*, \mathcal{M}_\Sigma \rangle &= \langle \mathcal{M}_A^* \otimes \mathcal{M}_B^*, \delta(\mathcal{M}_\Sigma) \rangle = \langle \mathcal{M}_A^* \otimes \mathcal{M}_B^*, \sum_{\Sigma = V \triangle U} \mathcal{M}_U \otimes \mathcal{M}_V \rangle \\ &= \sum_{\Sigma = V \triangle U} \langle \mathcal{M}_A^*, \mathcal{M}_U \rangle \langle \mathcal{M}_B^*, \mathcal{M}_V \rangle. \end{aligned}$$

This is 1 exactly when $\Sigma = B \triangle A$, otherwise it is 0. Thus it is equal to $\langle \mathcal{M}_{B \triangle A}^*, \mathcal{M}_\Sigma \rangle$, which proves the formula.

Note that the product Δ on tableaux turns T into a free monoid. This is proved similarly as in [13], where the similar result for the product \square is proved. The generators of this free monoid are the Δ -indecomposable tableaux.

Finally, we have the following criterion for the Δ -decomposability of a tableau (this criterion is however not very practical): let $r(T)$ be the row word associated to T , obtained by reading the entries of T in the reading order, that is from left to right and from top to bottom; then the tableau T is decomposable for Δ if, and only if, in the plactic class of $r(T)$ there is a permutation decomposable for Δ . Indeed, if T is obtained by dragging and dropping \bar{T}_2 on top of T_1 , then $r(T)$ is plactic-equivalent to $\overline{r(T_2)}r(T_1)$: this is contained in [13], p. 88 section e. (actually, for the product \square). The reciprocal is analogous: if in the plactic class of $r(T)$ there is a word decomposable for Δ , then T is decomposable for Δ . These observations follow by the well-known properties of the Robinson-Schensted correspondence and from the Eq. (2):

$$P(v \Delta u) = P(v) \Delta P(u).$$

7.3. A counter-example by Franco Saliola

Our basis \mathcal{M}_Σ was inspired by the construction of Aguiar and Sottile in [1]. They prove further that the structure constants for multiplication are positive (Theorem 4.1 in their article). This is

not true in the case of Poirier–Reutenauer Hopf algebra of tableaux, as shows a counter-example of Franco Saliola, that he kindly permitted us to reproduce here. Indeed, he computed that

$$\begin{aligned} \mathcal{M}_{P(123)} * \mathcal{M}_{P(123)} &= \mathcal{M}_{P(123456)} \\ &- \mathcal{M}_{P(241356)} - \mathcal{M}_{P(251346)} - \mathcal{M}_{P(261345)} - \mathcal{M}_{P(351236)} - \mathcal{M}_{P(361245)} - \mathcal{M}_{P(461235)} \\ &+ \mathcal{M}_{P(256134)} + \mathcal{M}_{P(346125)} + \mathcal{M}_{P(356124)} + 2\mathcal{M}_{P(456123)} \\ &+ 2\mathcal{M}_{P(362514)} - \mathcal{M}_{P(462513)} - \mathcal{M}_{P(543126)}. \end{aligned}$$

Acknowledgments

We thank Franco Saliola, who allowed us to include a counter-example arising from his computations. This work was partially supported by a grant of NSERC (Canada) for the second author, and was partially done during an invitation of him to the university La Sapienza in 2019.

References

- [1] M. Aguiar, F. Sottile, Structure of the Malvenuto-Reutenauer Hopf algebra of permutations, *Adv. Math.* 191 (2005) 225–275.
- [2] D. Blessenohl, M. Schocker, *Noncommutative Character Theory of the Symmetric Group*, Imperial College Press, 2005.
- [3] R. Cori, Indecomposable permutations, hypermaps and labeled Dyck paths, *J. Comb. Theory A* 116 (2009) 1326–1343.
- [4] G. Duchamp, F. Hivert, J.-C. Novelli, J.-Y. Thibon, Noncommutative symmetric functions VII: free quasi-symmetric functions revisited, *Ann. Comb.* 15 (2011) 655–673.
- [5] I.M. Gessel, Multipartite P -partitions and inner products of skew Schur functions, in: *Combinatorics and Algebra* (Boulder, Colo. 1983), in: *Contemporary Mathematics*, vol. 34, American Mathematical Society, Providence, RI, 1984, pp. 289–317.
- [6] <http://oeis.org>.
- [7] A. Jöllenbeck, Nichtkommutative Charaktertheorie der symmetrischen Gruppen, *Bayreuth. Math. Schr.* 56 (1999) 1–41.
- [8] G. Karaali, I. Senturia, M. Taşkın, A new partial order on SYT, preprint on arXiv.
- [9] A. Lascoux, M.-P. Schützenberger, Le monoïde plaxique, *Noncommutative structures in geometric Combinatorics*, Napoli, 1978, in: *Quaderni Della Ricerca Scientifica*, Vol. 109, CNR, Roma, 1981, pp. 129–156.¹
- [10] J.-L. Loday, M. Ronco, Order Structure on the Algebra of Permutations and of Planar Binary Trees, *J. Algebraic Combin.* 15 (2002) 253–270.
- [11] C. Malvenuto, C. Reutenauer, Duality between quasi-symmetric functions and the Solomon descent algebra, *J. Algebra* 177 (1995) 967–982.
- [12] A. Melnikov, On orbital variety closures of sl_n , I. Induced Duflo order, *J. Algebra* 271 (2004) 179–233.
- [13] S. Poirier, C. Reutenauer, Algèbres de Hopf de tableaux, *Ann. Sci. Math. Québec* 19 (1995) 79–90.
- [14] B. Sagan, *The Symmetric Group*, second ed., Springer, 2001.
- [15] L. Solomon, A mackey formula in the group ring of a Coxeter group, *J. Algebra* 41 (2) (1976) 255–264.
- [16] M. Taskin, Properties of four partial orders on standard Young tableaux, *J. Comb. Theory A* 113 (2006) 1092–1119.

¹ The collected papers of Schützenberger are available on the website of Jean Berstel.